

## **Sensitivity Analyses of Symptom-Based Progression-Free Survival as a Composite Endpoint in Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma: Results from the ALPINE Trial**

Jennifer R Brown<sup>1\*</sup>, Timothy Victor<sup>2,3</sup>, Nicole Lamanna<sup>4</sup>, Constantine Tam<sup>5</sup>, Lugui Qiu<sup>6</sup>, Rasika Korde<sup>3</sup>, Tommi Salmi<sup>7</sup>, Gisoo Barnes<sup>3</sup>, Mazyar Shadman<sup>8,9</sup>

<sup>1</sup>Dana-Farber Cancer Institute, Boston, MA, USA; <sup>2</sup>University of Pennsylvania, Philadelphia, PA, USA;

<sup>3</sup>BeOne Medicines USA, Inc., San Carlos, CA, USA; <sup>4</sup>Herbert Irving Comprehensive Cancer Center,

Columbia University, New York, NY, USA; <sup>5</sup>Alfred Health & Monash University, Melbourne, VIC,

Australia; <sup>6</sup>State Key Laboratory of Experimental Hematology, National Clinical Medical Research

Center for Blood Diseases, Institute of Hematology and Blood Diseases Hospital, Chinese Academy of

Medical Sciences and Peking Union Medical College, Tianjin, China; <sup>7</sup>BeOne Medicines Ltd,

Switzerland, GmbH, Basel, Switzerland; <sup>8</sup>Fred Hutchinson Cancer Center, Seattle, WA, USA;

<sup>9</sup>University of Washington, Seattle, WA, USA

**Presenting Author:** Dr. Jennifer Brown\*

**KEYWORDS:** Symptom-based progression-free survival, R/R CLL, zanubrutinib, Bruton tyrosine kinase inhibitor, sensitivity analysis

**BACKGROUND:** Symptom-based progression-free survival (S-PFS) is an emerging patient-centric composite endpoint integrating clinical disease progression with patient-reported outcomes (PROs). While PRO-PFS associations supporting S-PFS were demonstrated in the ALPINE trial (Brown et al. ASH 2025), their robustness under different death-handling assumptions is not established. The current analyses evaluated the robustness of previously reported PRO-PFS associations and zanubrutinib's treatment effect on PROs in chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL) under differing death-handling assumptions.

**METHODS:** Two post-hoc sensitivity analyses were conducted: (1) excluding all deaths regardless of cause (2) excluding non-disease-related deaths, as adjudicated by study investigator. Longitudinal PRO trajectories were assessed using the EORTC QLQ-C30, including 4 symptom domains (fatigue, insomnia, nausea/vomiting, pain) alongside GHS/QoL and physical functioning. A joint longitudinal-survival model was used; survival submodel results including treatment effects and PRO-PFS association parameters are reported.

**RESULTS:** Excluding all deaths (ibrutinib, n=228; zanubrutinib, n=245; [deaths excluded: ibrutinib n=76, zanubrutinib n=67]), zanubrutinib was associated with a significantly lower risk of disease progression for physical functioning versus ibrutinib (hazard ratio [HR] 0.60, 95% confidence interval [95%CI] 0.32–0.96;  $p=0.032$ ). Excluding non-disease-related deaths (ibrutinib, n=252; zanubrutinib, n=266; [deaths excluded: ibrutinib n=52, zanubrutinib n=46]), zanubrutinib was associated with a significantly lower risk of disease progression or death versus ibrutinib for GHS/QoL (HR 0.64, 95%CI 0.37–0.99;  $p=0.044$ ), nausea/vomiting (HR 0.65, 95%CI 0.38–0.99;  $p=0.045$ ), pain (HR 0.64, 95%CI 0.38–0.98;  $p=0.041$ ), and physical functioning (HR 0.61, 95%CI 0.34–0.95;  $p=0.027$ ). Excluding all deaths, worsening trajectories in fatigue (association parameter HR 1.13, 95%CI 1.05–1.30;  $p=0.007$ ) and insomnia (HR 1.12, 95%CI 1.02–1.31;  $p=0.009$ ) were significantly associated with increased risk of disease progression; both associations remained significant when excluding non-disease-related deaths, irrespective of treatment (fatigue: HR 1.12, 95%CI 1.03–1.28;  $p=0.004$ ; insomnia: HR 1.11, 95%CI 1.02–1.28;  $p=0.016$ ). Excluding non-disease-related deaths, higher physical functioning scores were significantly associated with lower risk of disease progression or disease-related death, irrespective of treatment (association parameter HR 0.96, 95%CI 0.92–1.00;  $p=0.031$ ).

**CONCLUSIONS:** Findings were consistent with previously reported PRO-PFS associations under both death-handling assumptions, supporting the robustness of zanubrutinib's patient-relevant treatment benefit versus ibrutinib and the credibility of S-PFS as a future endpoint in CLL/SLL trials.

**Sponsor:** BeOne Medicines, Ltd