

Direct medical costs of nasopharyngeal carcinoma in Indonesia: a healthcare payer perspective

Erna Kristin,¹ Cosphiadi Irawan,² Susanna Hilda Hutajulu,³ Yussy Afriani Dewi,⁴ Gregorius Ben Prajogi,⁵ Lucia Rizka Andalucia,⁶ Donni Hendrawan,⁷ Sudi Indrajaya,¹ Royasia Viki Ramadani,⁸ See-Hwee Yeo,⁸ Xiaowei Zhang,⁹ Junice Ng¹⁰

¹Faculty of Medicine, Public Health, and Nursing, Gadjah Mada University, Yogyakarta, Indonesia; ²Internal Medicine, Dr. Cipto Mangunkusumo Hospital, Universitas Indonesia, Jakarta, Indonesia; ³Division of Hematology and Medical Oncology, Department of Internal Medicine, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada/Dr Sardjito General Hospital, Yogyakarta, Indonesia; ⁴Department of Otorhinolaryngology-Head and Neck Surgery, Faculty of Medicine Padjadjaran University, Hasan Sadikin General Hospital, Bandung, Indonesia; ⁵Radiation Oncology, Dr. Cipto Mangunkusumo Hospital, Jakarta, Indonesia; ⁶Department of Pharmaceutical and Medical Devices, Ministry of Health, Indonesia; ⁷Research, Innovation and Development Department, BPJS Kesehatan, Indonesia; ⁸Real World Solutions, IQVIA Solutions Asia, Singapore, Republic of Singapore; ⁹BeOne Medicines, Inc., Global Statistics & Data Science, Shanghai, China; ¹⁰BeOne Medicines, Inc., Global HEOR, Singapore, Republic of Singapore

Background

- Context:** Nasopharyngeal carcinoma (NPC) represents a significant public health and economic burden, characterized by high mortality in Indonesia. A study conducted in Yogyakarta, Indonesia, reported a 35% overall survival rate at five years.¹ However, the specific costs it imposes on the healthcare system remain unclear.
- Study objective:** To estimate the direct medical costs of NPC from a public payer's perspective in Indonesia, using the national health insurance (JKN) database

Methods

- Study design:** Retrospective database study using information from the JKN database between January 1, 2018 and December 31, 2023
- About the database:** It covers 267 million Indonesians (~96% of the population) in 2023. It contains longitudinal data for demographics, diagnoses, treatment, healthcare resource utilization and costs. Disease diagnoses (primary and secondary diagnoses) are coded using the ICD-10, while treatment procedures are coded using the ICD-9-CM.
- About the healthcare payment mechanism:** This study captured two forms of hospital healthcare payment mechanisms, known as **INA-CBGs** (case-based groups; similar to diagnosis-related groups) and unbundling costs known as **Non-CBGs** tariffs.

Bundling costs (CBGs)

- Inpatient
 - Outpatient specialist
- These may include procedures, 7-day chemotherapy and non-cancer drugs, radiotherapy and surgery.

Unbundling costs (non-CBGs)

- Chemotherapy drugs beyond 7-days prescription
- Diagnostic and procedure
- Non-cancer drug
- Radiotherapy costs with special procedure


Hospital-based	Primary Care
Based on diagnosis group (CBGs)	Capitation based on catchment populations
Unbundling drugs and treatment/procedure (Non-CBGs)	Non-Capitation (fee-for-service)

CBGs, Case based groups

- Cohort:** NPC adult patients aged ≥18 years old with at least 2 medical visits related to NPC were identified from the JKN database between 2019 and 2022
 - Index date was defined as the first visit associated with NPC, as coded by ICD-10 'C11'
 - Baseline characteristics were assessed over a 12-month period before the index date
 - Patients were classified as treated for NPC if they received chemotherapy, radiotherapy, or surgery
- Analysis:** Results were summarized using descriptive statistics. Total costs represent sum of INA-CBGs and non-CBGs within 365 days after the index date. Costs in Indonesian Rupiah (IDR) were inflated using the Consumer Price Index in 2024 and converted to US dollars (US\$) (US\$ 1=IDR 15,881)

Results - Identified study population

- Incidence:** Among the 267 million patients with records in JKN, 23,072 patients were newly diagnosed with NPC between 2019 and 2022.
- Patients who received NPC-related treatment within 1 year after diagnosis:** 13,696 (59%)



- Did not receive treatment
- Chemotherapy only
- Chemotherapy and radiotherapy
- Radiotherapy only

Results

Demographic characteristics


Characteristic	Received NPC-related treatment (n=13,696)	Did not receive NPC-related treatment (n=9,376)	Total (N=23,072)
Male, n(%)	9,515 (69)	6,123 (65)	15,638 (68)
Age, mean (SD)	49.8 (12.2)	51.5 (13.7)	50.5 (12.9)
Residing in Java, n(%)	8,092 (59)	6,192 (66)	14,284 (62)
CCI≥5, n(%)	2,695 (20)	1,013 (11)	3,708 (16)
Subsidized, n(%)	4,987 (36)	4,048 (43)	9,035 (39)

*Charlson Comorbidity Index

Direct medical cost (in USD) for the cohort

Direct medical cost	Received NPC-related treatment (n=13,696)	Did not receive NPC-related treatment (n=9,376)	Total (N=23,072)
Mean (SD)	3,227 (2,597)	499 (910)	2,118 (2,477)
Median (IQR)	2,874 (3,308)	230 (482)	1,107 (3,198)

Direct medical cost (in USD) by treatment modality



Conclusion

- This is the **first nationwide study** to estimate the direct medical cost of NPC based upon the public payer to help inform decision-makers
- The total cost in 2022 was estimated at US\$ 14.8 million, or **5% of the JKN cancer expenditure**
- Despite the significant spending on managing this disease, mortality rates remain high, highlighting a **critical gap in the effectiveness of current treatment options**. In particular, radiotherapy utilization rate was low suggesting late diagnosis and poor access to curative treatment

Reference

- Hutajulu SH et al. PLoS One. 2021;16(2):e0246638
- BPJS Kesehatan, Program Management Report & Financial Report 2022

Acknowledgments

- This study was sponsored by BeOne Medicines, Ltd. The analysis was provided by Real World Solutions, IQVIA Solutions Asia; editorial and submission support was provided by Paraxel and was funded by BeOne Medicines
- We would like to thank *Badan Penyelenggara Jaminan Sosial Kesehatan* for providing the data